

Heat Stability of TREHALOSE SG

Material

TREHALOSE SG Lot No. 4D04

Preparation of test solutions

TREHALOSE SG (dihydrate crystal) aqueous solutions were prepared at concentrations of 33, 40 and 50 g/L.

Heating conditions of solutions and evaluation of Trehalose content

Forty (40) mL each of the Trehalose solutions were poured into glass containers and heated at 122°C for 0, 20, 40, 60 or 120 min.



The solutions were analyzed by HPLC and the Trehalose content (%) was calculated by the Area normalization method.

Results

No changes in Trehalose content were observed even when the solutions were heated at 122°C for 120 min.

Trehalose content (%) before and after heating

Heating condition	0 min	122°C 20 min	122°C 40 min	122°C 60 min	122°C 120 min
Trehalose 33 g/L	99.96	99.95	99.96	99.95	99.96
Trehalose 40 g/L	99.96	99.96	99.96	99.95	99.96
Trehalose 50 g/L	99.96	99.96	99.96	99.96	99.95



Chromatograms of TREHALOSE SG solution (50 g/L) heated at 122°C for 0 min left, and 120 min right.

MANUFACTURER : **Nagase Viita Co., Ltd.**
CONTACT : **Nagase & Co., Ltd.**

Life & Healthcare Products Department
E-mail: dnfct@ex.nagase.co.jp

⚠ The information provided herein is intended only for reference purposes. It is the customer's responsibility to determine that the ingredients meet all legal requirements in the country where they are used, and that they do not infringe on any third party patents. Unauthorized reproduction of this brochure is prohibited.